

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PW 279062
(M#)

Invention: NUCLEOTIDE SEQUENCES ENCODING A SENSOR KINASE, *citA*, FROM
CORYNEBACTERIUM GLUTAMICUM

Inventor (s): Bettina MÖCKEL
Mike FARWICK
Thomas HERMANN
Achim MARX
Walter PFEFFERLE



For correspondence Address



00909

Pillsbury Winthrop LLP

This is a:

- ☐ Provisional Application
- ☐ Regular Utility Application
- ☐ Continuing Application
 - ☒ The contents of the parent are incorporated by reference
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification
 - Sub. Spec Filed _____
 - in App. No. _____
- ☒ Marked up Specification re
 - Sub. Spec. filed March 6, 2003
 - In App. No. 09/804,060

SPECIFICATION



[New nucleotide sequences which code for the citA gene]
Nucleotide Sequences Encoding a Sensor Kinase,
citA, from Corynebacterium Glutamicum

The invention provides nucleotide sequences from coryneform
5 bacteria which code for the citA gene and a process for the
fermentative preparation of amino acids, in particular L-
lysine, by attenuation of the citA gene. The citA gene
codes for the sensor kinase Cit A of a two-component
system.

10 [Prior art] Background

L-Amino acids, in particular L-lysine, are used in human
medicine and in the pharmaceuticals industry, in the
foodstuffs industry and very particularly in animal
nutrition.

15 It is known that amino acids are prepared by fermentation
from strains of coryneform bacteria, in particular
Corynebacterium glutamicum. Because of their great
importance, work is constantly being undertaken to improve
the preparation processes. Improvements to the process can
20 relate to fermentation measures, such as, for example,
stirring and supply of oxygen, or the composition of the
nutrient media, such as, for example, the sugar
concentration during the fermentation, or the working up to
the product form by, for example, ion exchange
25 chromatography, or the intrinsic output properties of the
microorganism itself.

Methods of mutagenesis, selection and mutant selection are
used to improve the output properties of these
microorganisms. Strains which are resistant to
30 antimetabolites or are auxotrophic for metabolites of
regulatory importance and which produce amino acids are
obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acids.

Object of the Invention

- 5 The inventors had the object of providing new measures for improved fermentative preparation of amino acids, in particular L-lysine.

Brief Description of the Invention

- If L-lysine or lysine are mentioned in the following, this
10 also means the salts, such as e.g., lysine monohydrochloride or lysine sulfate.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the *citA* gene chosen from the group
15 consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- 20 b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- c) polynucleotide which is complementary to the
25 polynucleotides of a) or b), and
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of sensor
30 kinase CitA.

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1 or
- 5 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequences complementary to sequences (i) or (ii),
10 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides:

a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as
15 shown in SEQ ID No.1;

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing parts of the polynucleotide according
20 to the invention, but at least 15 successive nucleotides of the sequence claimed

and coryneform bacteria in which the citA gene is attenuated, in particular by an insertion or deletion.

The invention also provides polynucleotides which
25 substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide
30 according to the invention or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

Polynucleotides comprising the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for
5 CitA protein or to isolate those nucleic acids or polynucleotides or genes which have a high similarity with the sequence of the citA gene.

Polynucleotides comprising the sequences according to the invention are furthermore suitable as primers with the aid
10 of which DNA of genes which code for the CitA protein can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides which serve as probes or primers comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides.
15 Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

Detailed Description of the Invention

The following figure is attached:

Figure 1: Map of the plasmid pCR2.1citAint.

20 The abbreviations and designations used have the following meaning.

<u>KmR:</u>	<u>Kanamycin resistance gene</u>
<u>EcoRI:</u>	<u>Cleavage site of the restriction enzyme</u> <u>EcoRI</u>
<u>HindIII:</u>	<u>Cleavage site of the restriction enzyme</u> <u>HindIII</u>
<u>BamHI:</u>	<u>Cleavage site of the restriction enzyme</u> <u>BamHI</u>
<u>citAint:</u>	<u>Internal fragment of the citA gene</u>

ColE1: Replication origin of the plasmid ColE1

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these
5 to be non-modified RNA or DNA or modified RNA or DNA.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70%, preferably at least 80% and in particular at least 90% to
10 95% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

15 The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of the CitA protein and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the
20 polypeptide according to SEQ ID No. 2 and have the activity mentioned.

The invention moreover relates to a process for the fermentative preparation of amino acids, in particular L-lysine, using coryneform bacteria which in particular
25 already produce amino acids and in which the nucleotide sequences which code for the citA gene are attenuated, in particular eliminated or expressed at a low level.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of
30 one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak

promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

5 The microorganisms which the present invention provides can prepare amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of
10 the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in
15 particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

Corynebacterium glutamicum ATCC13032
Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
20 *Corynebacterium melassecola* ATCC17965
Corynebacterium thermoaminogenes FERM BP-1539
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869 and
Brevibacterium divaricatum ATCC14020

25 or L-amino acid-producing mutants or strains prepared therefrom, such as, for example, the L-lysine-producing strains

Corynebacterium glutamicum FERM-P 1709
Brevibacterium flavum FERM-P 1708
30 *Brevibacterium lactofermentum* FERM-P 1712
Corynebacterium glutamicum FERM-P 6463
Corynebacterium glutamicum FERM-P 6464
Corynebacterium glutamicum DM58-1
Corynebacterium glutamicum DG52-5

Corynebacterium glutamicum DSM 5715 and
Corynebacterium glutamicum DSM 12866

The inventors have succeeded in isolating the new *citA* gene of *C. glutamicum* which codes for the CitA protein and which
5 is a sensor kinase of a two-component system.

To isolate the *citA* gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks
10 and handbooks. The textbook by Winnacker: [Gene und Klone, Eine Einführung in die Gentechnologie [] Genes and Clones, An Introduction to Genetic Engineering []] (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: Molecular Cloning, A Laboratory Manual
15 (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ -vectors by Kohara et al. (Cell 50, 495-508 (1987)). Bathe et al. (Molecular and General Genetics, 252:255-265, 1996)
20 describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research
25 16:1563-1575). Börmann et al. (Molecular Microbiology 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pHC79 (Hohn and Collins, 1980, Gene 11, 291-298).

To prepare a gene library of *C. glutamicum* in *E. coli* it is
30 also possible to use plasmids such as pBR322 (Bolivar, 1979, Life Sciences, 25, 807-818) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective, such as, for example, the
35 strain DH5 α (Jeffrey H. Miller: "A Short Course in Bacterial Genetics, A Laboratory Manual and Handbook for

Escherichia coli and Related Bacteria", Cold Spring
[Harbour [sic]] Harbor Laboratory Press, 1992).

The long DNA fragments cloned with the aid of cosmids or
other λ -vectors can than be subcloned in turn into the
5 usual vectors suitable for DNA sequencing.

Methods of DNA sequencing are described, inter alia, by
Sanger et al. (Proceedings of the National Academy of
Sciences of the United States of America USA, 74:5463-5467,
1977).

10 The resulting DNA sequences can then be investigated with
known algorithms or sequence analysis programs, such as
e.g., that of Staden (Nucleic Acids Research 14, 217-
232(1986)), that of Marck (Nucleic Acids Research 16, 1829-
1836 (1988)) or the GCG program of Butler (Methods of
15 Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the
citA gene and which, as SEQ ID No. 1, is a constituent of
the present invention has been found in this way. The
amino acid sequence of the corresponding protein has
20 furthermore been derived from the present DNA sequence by
the methods described above. The resulting amino acid
sequence of the *citA* gene product is shown in SEQ ID No. 2.

Coding DNA sequences which result from SEQ ID No. 1 by the
degeneracy of the genetic code are also a constituent of
25 the invention. In the same way, DNA sequences which
hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a
constituent of the invention. Conservative amino acid
exchanges, such as e.g., exchange of glycine for alanine or
of aspartic acid for glutamic acid in proteins, are
30 furthermore known among experts as "sense mutations" which
do not lead to a fundamental change in the activity of the
protein, i.e., are of neutral function. It is furthermore
known that changes on the N and/or C terminus of a protein
cannot substantially impair or can even stabilize the

function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 5 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

10 Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

15 Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal 20 of Systematic Bacteriology 41: 255-260 (1991)). The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i.e., the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that 25 the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps 30 (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996). A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the 35 sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This

can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50 to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g., DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: [OligonukleotidesA: [sic] a] Oligonucleotides: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

In the work on the present invention, it has been found that coryneform bacteria produce amino acids, in particular L-lysine, in an improved manner after attenuation of the *citA* gene.

To achieve an attenuation, either the expression of the *citA* gene or the catalytic properties of the enzyme protein can be reduced or eliminated. The two measures can optionally be combined.

The reduction in gene expression can take place by suitable culturing or by genetic modification (mutation) of the signal structures of gene expression. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information on this e.g., in the patent

application WO 96/15246, in Boyd and Murphy (Journal of Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss (Nucleic Acids Research 26: 3548 (1998)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)),
5 in Pátek et al. (Microbiology 142: 1297 (1996)), Vasicova et al. (Journal of Bacteriology 181: 6188 (1999)) and in known textbooks of genetics and molecular biology, such as e.g., the textbook by Knippers ("[Molekulare Genetik
[Molecular Genetics]]," 6th edition, Georg Thieme Verlag,
10 Stuttgart, Germany, 1995) or that by Winnacker ("[Gene und Klone [Genes and Clones]]," VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the
15 prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("[Die Threonindehydratase aus Corynebacterium glutamicum:
20 Aufhebung der allosterischen Regulation und Struktur des Enzyms [Threonine dehydratase from Corynebacterium glutamicum: Cancelling the allosteric regulation and structure of the enzyme]]," Reports from the Jülich Research Centre, Jül-2906, ISSN09442952, Jülich, Germany,
25 1994). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g., that by Hagemann ("[Allgemeine Genetik [General Genetics]]," Gustav Fischer Verlag, Stuttgart, 1986).

Possible mutations are transitions, transversions,
30 insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, missense mutations or nonsense mutations are referred to. Insertions or deletions of at least one base pair (bp) in a gene lead to frame shift mutations, as a consequence of
35 which incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons

typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and molecular biology, such as e.g., the textbook by Knippers
5 ("[Molekulare Genetik []Molecular Genetics[]]," 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("[Gene und Klone []Genes and Clones[]]," VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("[Allgemeine Genetik []General Genetics[]]",
10 Gustav Fischer Verlag, Stuttgart, 1986).

A common method of mutating genes of *C. glutamicum* is the method of gene disruption and gene replacement described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the
15 coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-
20 73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US Patent 5,487,993), pCR®Blunt (Invitrogen,
25 Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516). The plasmid vector which contains the central part of the coding region of the gene is then transferred into the desired strain of
30 *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and
35 Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS

Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross-over" event, the coding region of the gene in question is interrupted by the vector sequence and two incomplete alleles are
5 obtained, one lacking the 3' end and one lacking the 5' end. This method has been used, for example, by Fitzpatrick et al. (Applied Microbiology and Biotechnology 42, 575-580 (1994)) to eliminate the *recA* gene of *C. glutamicum*.

10 In the method of gene replacement, a mutation, such as e.g., a deletion, insertion or base exchange, is established in vitro in the gene of interest. The allele prepared is in turn cloned in a vector which is not replicative for *C. glutamicum* and this is then transferred
15 into the desired host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross-over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the
20 incorporation of the mutation or of the allele is achieved. This method was used, for example, by Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to eliminate the *pyc* gene of *C. glutamicum* by a deletion.

A deletion, insertion or a base exchange can be
25 incorporated into the *citA* gene in this manner.

In addition, it may be advantageous for the production of L-amino acids, in particular L-lysine, to enhance, in particular to over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of
30 anaplerosis, of the pentose phosphate cycle or of amino acid export and optionally regulatory proteins, in addition to attenuation of the *citA* gene.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more
35 enzymes (proteins) in a microorganism which are coded by

the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding enzyme (protein) having a high activity, and optionally
5 combining these measures.

Thus, for example, for the preparation of L-lysine, at the same time one or more of the genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate
10 synthase (EP-B 0 197 335),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the zwf gene which codes for glucose 6-phosphate
15 dehydrogenase (JP-A-09224661),
- the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
- the lysE gene which codes for lysine export (DE-A-195 48 222)
- 20 • the lysC gene which codes for a feed back resistant aspartate kinase (EP-B-0387527; EP-A-0699759; WO 00/63388), or
- the zwal gene which codes for the Zwal protein (DE: 199 59 328.0, DSM 13115)

25 can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of amino acids, in particular L-lysine, in addition to the attenuation of the citA gene, at the same time for one or more of the genes chosen from the group consisting of

- the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM 13047),
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),
- 5 • the poxB gene which codes for pyruvate oxidase (DE:1995 1975.7, DSM 13114)
- the zwa2 gene which codes for the Zwa2 protein (DE: 199 59,327.2, DSM 13113)

to be attenuated.

- 10 In addition to attenuation of the citA gene it may furthermore be advantageous, for the production of amino acids, in particular L-lysine, to eliminate undesirable side reactions, (Nakayama: "Breeding of Amino Acid Producing Microorganisms," in: Overproduction of Microbial
- 15 Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch

20 culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of L-amino acids, in particular L-lysine. A summary of known culture methods is described in the textbook by Chmiel ([Bioprozesstechnik 1. Einführung in die

25 Bioverfahrenstechnik []Bioprocess Technology 1.

Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas ([Bioreaktoren und periphere Einrichtungen []Bioreactors and Peripheral Equipment[]] (Vieweg Verlag,

30 Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained

in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e.g., glucose, sucrose, lactose, fructose, maltose, 5 molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as, for example, palmitic acid, stearic acid and linoleic acid, alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, 10 for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep 15 liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

20 Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron 25 sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can 30 be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed 35 in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol

esters, can be employed to control the development of foam. Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic
5 conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is
10 usually reached within 10 hours to 160 hours.

Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by anion exchange
15 chromatography with subsequent ninhydrin derivatization, or it can be carried out by reversed phase HPLC, for example as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The invention furthermore relates to a process for the
20 fermentative preparation of an amino acid chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-
25 arginine, in particular L-lysine, using coryneform bacteria which in particular already produce one or more of the amino acids mentioned.

The following microorganism was deposited on 23.01.2001 as a pure culture at the [Deutsche Sammlung für Mikrorganismen
30 [sic] und Zellkulturen (DSMZ =] German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany[]] in accordance with the Budapest Treaty:

- Escherichia coli strain Top10/pCR2.1citAint as DSM 13998.

The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual, 1989, Cold Spring [Harbour [sic]] Harbor Laboratory Press, Cold Spring Harbor, NY, USA). Methods for transformation of Escherichia coli are also described in this handbook.

- 10 The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

ExamplesExample 1

Preparation of a genomic cosmid gene library from *C. glutamicum* ATCC 13032

- 5 Chromosomal DNA from *C. glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were
- 10 dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987), Proceedings of the National Academy of Sciences, USA 84:2160-2164), obtained
- 15 from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vektor Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline
- 20 phosphatase.

- The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032
- 25 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no. 27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product
- 30 Description Gigapack II XL Packing Extract, Code no. 200217).

For infection of the *E. coli* strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage

suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 5 1955, Virology, 1:190) + 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 2

Isolation and sequencing of the *citA* gene

10 The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product 15 Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid 20 fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description 25 Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out 30 as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol 35 Letters, 123:343-7) into the *E. coli* strain DH5αMCR (Grant,

1990, Proceedings of the National Academy of Sciences, U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 µg/ml zeocin.

The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academies of Sciences, U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analyses were carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402) against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1653 bp, which was called the *citA* gene. The *citA* gene codes for a polypeptide of [of [sic]] 551 amino acids.

Example 3

Preparation of an integration vector for integration
mutagenesis of the *citA* gene

From the strain ATCC 13032, chromosomal DNA was isolated by
5 the method of Eikmanns et al. (Microbiology 140: 1817 -
1828 (1994)). On the basis of the sequence of the *citA*
gene known for *C. glutamicum* from example 2, the following
oligonucleotides were chosen for the polymerase chain
reaction (see SEQ ID No. 4 and SEQ ID No. 5):

10 *citA*-int1:

5` TTC CAG TCG GTG AGG TCA GT 3`

citA-int2:

5` GTA CGA TCG CGG ATG GTT AC 3`

The primers shown were synthesized by MWG Biotech
15 (Ebersberg, Germany) and the PCR reaction was carried out
by the standard PCR method of Innis et al. (PCR protocols.
A guide to methods and applications, 1990, Academic Press)
with the Taq-polymerase from Boehringer Mannheim (Germany,
Product Description Taq DNA polymerase, Product No. 1 146
20 165). With the aid of the polymerase chain reaction, the
primers allow amplification of an internal fragment of the
citA gene 480 bp in size. The product amplified in this
way was tested electrophoretically in a 0.8% agarose gel.

The amplified DNA fragment (see SEQ ID No. 3) was ligated
25 with the TOPO TA Cloning Kit from Invitrogen Corporation
(Carlsbad, CA, USA; Catalogue Number K4500-01) in the
vector pCR2.1-TOPO (Mead et al. (1991) Bio/Technology
9:657-663).

The *E. coli* strain TOP10 was then electroporated with the
30 ligation batch (Hanahan, In: DNA cloning. A practical
approach. Vol. I. IRL-Press, Oxford, Washington DC, USA,
1985). Selection for plasmid-carrying cells was made by
plating out the transformation batch on LB agar (Sambrook
et al., Molecular cloning: a laboratory manual. 2nd Ed.

Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1citAint and is shown in figure 1.

The following microorganism was deposited as a pure culture on 23.01.2001 at the [Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ =] German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany[] in accordance with the Budapest Treaty:

- *Escherichia coli* Top10/pCR2.1citAint as DSM 13998

15 Example 4

Integration mutagenesis of the *citA* gene in the strains DSM 5715 and FERM-BP 1763

The vector pCR2.1citAint mentioned in example 3 was electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) into the strains *Corynebacterium glutamicum* DSM 5715 and *Brevibacterium lactofermentum* FERM-BP 1763. The strain DSM 5715 is an AEC-resistant lysine producer (EP-B-0435132), and the strain FERM-BP 1763 is a mycophenolic acid-resistant valine producer (US-A-5,188,948). The vector pCR2.1citAint cannot replicate independently in DSM5715 and FERM-BP 1763 and is retained in the cell only if it has integrated into the chromosome of DSM 5715 or FERM-BP 1763. Selection of clones with pCR2.1citAint integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin.

For detection of the integration, the citAint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of in each case a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes EcoRI, BamHI and HindIII. The fragments formed were separated by means of agarose gel electrophoresis and hybridized at 68°C with the Dig [hybridization [sic]] hybridization kit from Boehringer. The plasmid pCR2.1citAint mentioned in example 3 had been inserted into the chromosome of DSM5715 and into the chromosome of FERM-BP 1763 within the chromosomal cita gene. The strains were called DSM5715::pCR2.1citAint and FERM-BP 1763::pCR2.1citAint.

Example 5

Preparation of lysine

The C.glutamicum strain DSM5715::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l) [sic]) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
Bacto-Peptide	10 g/l
Bacto-Yeast extract	10 g/l
Glucose (autoclaved separately)	2% (w/v)

The pH is brought to pH 7.4

Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of the main culture is 0.1 OD. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
$[(\text{NH}_4)_2\text{SO}_4]$ [sic] <u>$(\text{NH}_4)_2\text{SO}_4$</u>	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 * 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 * 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 * 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 * \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l

Leucine (sterile-filtered) 0.1 g/l

CaCO₃ 25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, and the CaCO₃ autoclaved in the dry state was added.

- 5 Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD was determined at a measurement
10 wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

- 15 The result of the experiment is shown in table 1.

Table 1

Strain	OD(660)	Lysine HCl g/l
DSM5715	7.5	13.3
DSM5715::pCR2.1citAint	7.4	14.4

Example 6

Preparation of valine

- 20 The B. lactofermentum strain FERM-BP 1763::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium

suitable for the production of valine and the valine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with 5 kanamycin (25 mg/l) ~~[[sic]]~~ for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture. Kanamycin (25 mg/l) was added to this. The preculture was incubated 10 for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1 OD. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
$[(\text{NH}_4)_2\text{SO}_4]$ [[sic]] <u>$(\text{NH}_4)_2\text{SO}_4$</u>	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 * 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 * 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 * 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 * \text{H}_2\text{O}$	5.0 mg/l
Isoleucine (sterile-filtered)	0.1 g/l
Methionine (sterile-filtered)	0.1 g/l

Thiamine * HCl (sterile-filtered) 0.2 mg/l
Leucine (sterile-filtered) 0.1 g/l
CaCO₃ 25 g/l

The CSL (corn steep liquor), MOPS (morpholinopropanesulfonic acid) and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, 5 and the CaCO₃ autoclaved in the dry state was added.

Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

10 After 48 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of valine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography 15 and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 2.

Table 2

Strain	OD(660)	Valine HCl g/l
FERM-BP 1763	12.1	7.5
FERM-BP 1763::pCR2.1citAint	13.5	10.8

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG

5 <120> New nucleotide sequences which code for the citA gene

<130> 000169 BT

<140>

10 <141>

<160> 5

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 2055

<212> DNA

<213> Corynebacterium glutamicum

20

<220>

<221> CDS

<222> (201) .. (1853)

<223> citA gene

25

<400> 1

tttctgtgtt tctcgaaact ttgagatccc gagtggtctg tgttgcttgt gggagtataa 60

gggtggcgcgt gtcacgcaca gaagtgtttg gtgcattgcc tgaggtagtg cgcaaaataa 120

30

gactttttgtg cattatgatc agaattgttg gcctgggact tcgcttcacg ctctgctgat 180

aatcgccccc gggggtagac atg tct gtt ggt gga tcc gac tgg aaa aac ttc 233

35

Met Ser Val Gly Gly Ser Asp Trp Lys Asn Phe

1

5

10

aag gag gtg gac atc att cgc ttt gct acc cga ata ctg gtg att caa 281

Lys Glu Val Asp Ile Ile Arg Phe Ala Thr Arg Ile Leu Val Ile Gln

15

20

25

40

gtg gct acc gtc gcg ttg gtg gta gct att tgc acc gga att ttc gca 329

Val Ala Thr Val Ala Leu Val Val Ala Ile Cys Thr Gly Ile Phe Ala

30

35

40

45

gtt ttg atg atg gat cag atg aaa act gag gcc gag cac aca gcg ctg 377

Val Leu Met Met Asp Gln Met Lys Thr Glu Ala Glu His Thr Ala Leu

45

50

55

tcc atc gga cgt tcg gtg gca tcc aac ccg cag atc cgc gag gaa gta 425

50

Ser Ile Gly Arg Ser Val Ala Ser Asn Pro Gln Ile Arg Glu Glu Val

60

65

70

75

gcg ctt gat act caa aca gga gca aac cca tcg gcc gaa gaa tta gcc 473

55

Ala Leu Asp Thr Gln Thr Gly Ala Asn Pro Ser Ala Glu Glu Leu Ala

80

85

90

gat gga gat atc caa gcg gtt gca cag gcg gcc aat gaa cgc act gga 521

Asp Gly Asp Ile Gln Ala Val Ala Gln Ala Ala Asn Glu Arg Thr Gly

95

100

105

60

	gct ttg ttt gtc gtt atc act gac ggt tta ggt atc cgc ctg tcc cac	569
	Ala Leu Phe Val Val Ile Thr Asp Gly Leu Gly Ile Arg Leu Ser His	
	110 115 120	
5	cca gat gag gaa cgt ctg ggg gag cag gtg agc act agc ttt gag gct	617
	Pro Asp Glu Glu Arg Leu Gly Glu Gln Val Ser Thr Ser Phe Glu Ala	
	125 130 135	
10	gcc atg cgg ggt gaa gaa acc atg gcg tgg gag act ggg acc ctc ggt	665
	Ala Met Arg Gly Glu Glu Thr Met Ala Trp Glu Thr Gly Thr Leu Gly	
	140 145 150 155	
15	gcg tcc gcg cga gca aaa gtg cct atc ttt gcg ccg gat tct agt gtt	713
	Ala Ser Ala Arg Ala Lys Val Pro Ile Phe Ala Pro Asp Ser Ser Val	
	160 165 170	
20	cca gtc ggt gag gtc agt gtt ggg ttt gag cga gac agt gtg tat tcc	761
	Pro Val Gly Glu Val Ser Val Gly Phe Glu Arg Asp Ser Val Tyr Ser	
	175 180 185	
25	cgc ctg ccc atg ttc ctc gcc gcc ctt gct ctt att tct gtg ttg gga	809
	Arg Leu Pro Met Phe Leu Ala Ala Leu Ala Leu Ile Ser Val Leu Gly	
	190 195 200	
30	atc ctt atc ggc gtg ggt gta gcc atg ggc atg cga cgc cgt tgg gaa	857
	Ile Leu Ile Gly Val Gly Val Ala Met Gly Met Arg Arg Arg Trp Glu	
	205 210 215	
35	cgc gtg acc ttg ggt ttg cag ccg gag gag cta gtg acc ctt gtg caa	905
	Arg Val Thr Leu Gly Leu Gln Pro Glu Glu Leu Val Thr Leu Val Gln	
	220 225 230 235	
40	aat cag act gca gtc atc gat ggc att gat gag ggc gtg ctg gcg ctg	953
	Asn Gln Thr Ala Val Ile Asp Gly Ile Asp Glu Gly Val Leu Ala Leu	
	240 245 250	
45	agc cca aac gga aca att ggg gtg cat aat gag cag gcg caa tcc atg	1001
	Ser Pro Asn Gly Thr Ile Gly Val His Asn Glu Gln Ala Gln Ser Met	
	255 260 265	
50	att ggt gca ggt cct atg agt ggc agg acg ttg aaa gaa cta ggg ctt	1049
	Ile Gly Ala Gly Pro Met Ser Gly Arg Thr Leu Lys Glu Leu Gly Leu	
	270 275 280	
55	gac ctg ggt ctt gat ggc gtt gta ttg cat ggt cag cat ccg gaa acc	1097
	Asp Leu Gly Leu Asp Gly Val Val Leu His Gly Gln His Pro Glu Thr	
	285 290 295	
60	gtt gcc cat aac ggc agg atc ctc tat ctg gat ttc cac ccc gtg cgc	1145
	Val Ala His Asn Gly Arg Ile Leu Tyr Leu Asp Phe His Pro Val Arg	
	300 305 310 315	
65	cgt ggg gat caa gat tta ggc tac gtg gta acc atc cgc gat cgt acc	1193
	Arg Gly Asp Gln Asp Leu Gly Tyr Val Val Thr Ile Arg Asp Arg Thr	
	320 325 330	
70	gac atc att gaa ctc agt gaa cgc ctc gac tct gtg cgc acc atg acc	1241
	Asp Ile Ile Glu Leu Ser Glu Arg Leu Asp Ser Val Arg Thr Met Thr	
	335 340 345	

cac gca ctc cgc gcc cag cgc cac gag ttt gcc aac cgc atc cac acc 1289
 His Ala Leu Arg Ala Gln Arg His Glu Phe Ala Asn Arg Ile His Thr
 350 355 360

5 gca aca ggg ctt atc gac gcc ggc cgc gtc cac gac gcg gca gag ttt 1337
 Ala Thr Gly Leu Ile Asp Ala Gly Arg Val His Asp Ala Ala Glu Phe
 365 370 375

10 cta ggc gat ata tcc cgc aac ggg gga cag tca cat cca ttg atc gga 1385
 Leu Gly Asp Ile Ser Arg Asn Gly Gly Gln Ser His Pro Leu Ile Gly
 380 385 390 395

15 tca gcg cac ctc aat gaa gca ttt ttg agc tca ttt tta agt act gct 1433
 Ser Ala His Leu Asn Glu Ala Phe Leu Ser Ser Phe Leu Ser Thr Ala
 400 405 410

20 tct att tcg gca tct gaa aag ggc gtt agt ctg cgc atc aac tct gac 1481
 Ser Ile Ser Ala Ser Glu Lys Gly Val Ser Leu Arg Ile Asn Ser Asp
 415 420 425

acg ctc atc ctt ggc act gtt aaa gat cca gaa gat gta gca acc att 1529
 Thr Leu Ile Leu Gly Thr Val Lys Asp Pro Glu Asp Val Ala Thr Ile
 430 435 440

25 ttg ggt aat tta atc aac aat gcc atc gac gcc gcg gtg gca ggt gaa 1577
 Leu Gly Asn Leu Ile Asn Asn Ala Ile Asp Ala Ala Val Ala Gly Glu
 445 450 455

30 gcc cca cgg tgg att gag ctt acg ttg atg gat gat gcc gat acg ctg 1625
 Ala Pro Arg Trp Ile Glu Leu Thr Leu Met Asp Asp Ala Asp Thr Leu
 460 465 470 475

35 gtc att tct gtt gca gat tct ggt cct gga atc cca gag ggc gtg gat 1673
 Val Ile Ser Val Ala Asp Ser Gly Pro Gly Ile Pro Glu Gly Val Asp
 480 485 490

40 gta ttt gcc aca gcc acc cag ata gga gac tct gaa gat aat gaa cgc 1721
 Val Phe Ala Thr Ala Thr Gln Ile Gly Asp Ser Glu Asp Asn Glu Arg
 495 500 505

acc cac ggg cat ggc att ggt cta aaa ctg tgc cgg gct ttg gct aga 1769
 Thr His Gly His Gly Ile Gly Leu Lys Leu Cys Arg Ala Leu Ala Arg
 510 515 520

45 tca cat ggt ggc gat gtc tgg gtg att gat aga gga acc gaa gat ggc 1817
 Ser His Gly Gly Asp Val Trp Val Ile Asp Arg Gly Thr Glu Asp Gly
 525 530 535

50 gct gta ttt gga gtg aaa cta ccg gga gta atg gag taatggatca 1863
 Ala Val Phe Gly Val Lys Leu Pro Gly Val Met Glu
 540 545 550

aacacttaaa gtttttagtaa ttgatgatga tttccgcgtc gccggcattc acgcctccat 1923

55 cgttgatgcg tcccctggat tttcggtggt cggtaccgcg cgtaccctcg cagaggcaaa 1983

aaccctgate gccacatttt ccccggtatct cctacttggt gatgtctacc tccccgacgg 2043

60 cgatggcatt ga 2055

<210> 2
 <211> 551
 <212> PRT
 5 <213> Corynebacterium glutamicum

<400> 2
 Met Ser Val Gly Gly Ser Asp Trp Lys Asn Phe Lys Glu Val Asp Ile
 1 5 10 15
 10 Ile Arg Phe Ala Thr Arg Ile Leu Val Ile Gln Val Ala Thr Val Ala
 20 25 30
 15 Leu Val Val Ala Ile Cys Thr Gly Ile Phe Ala Val Leu Met Met Asp
 35 40 45
 Gln Met Lys Thr Glu Ala Glu His Thr Ala Leu Ser Ile Gly Arg Ser
 50 55 60
 20 Val Ala Ser Asn Pro Gln Ile Arg Glu Glu Val Ala Leu Asp Thr Gln
 65 70 75 80
 Thr Gly Ala Asn Pro Ser Ala Glu Glu Leu Ala Asp Gly Asp Ile Gln
 85 90 95
 25 Ala Val Ala Gln Ala Ala Asn Glu Arg Thr Gly Ala Leu Phe Val Val
 100 105 110
 30 Ile Thr Asp Gly Leu Gly Ile Arg Leu Ser His Pro Asp Glu Glu Arg
 115 120 125
 Leu Gly Glu Gln Val Ser Thr Ser Phe Glu Ala Ala Met Arg Gly Glu
 130 135 140
 35 Glu Thr Met Ala Trp Glu Thr Gly Thr Leu Gly Ala Ser Ala Arg Ala
 145 150 155 160
 Lys Val Pro Ile Phe Ala Pro Asp Ser Ser Val Pro Val Gly Glu Val
 165 170 175
 40 Ser Val Gly Phe Glu Arg Asp Ser Val Tyr Ser Arg Leu Pro Met Phe
 180 185 190
 45 Leu Ala Ala Leu Ala Leu Ile Ser Val Leu Gly Ile Leu Ile Gly Val
 195 200 205
 Gly Val Ala Met Gly Met Arg Arg Arg Trp Glu Arg Val Thr Leu Gly
 210 215 220
 50 Leu Gln Pro Glu Glu Leu Val Thr Leu Val Gln Asn Gln Thr Ala Val
 225 230 235 240
 Ile Asp Gly Ile Asp Glu Gly Val Leu Ala Leu Ser Pro Asn Gly Thr
 245 250 255
 55 Ile Gly Val His Asn Glu Gln Ala Gln Ser Met Ile Gly Ala Gly Pro
 260 265 270
 60 Met Ser Gly Arg Thr Leu Lys Glu Leu Gly Leu Asp Leu Gly Leu Asp
 275 280 285

Gly Val Val Leu His Gly Gln His Pro Glu Thr Val Ala His Asn Gly
 290 295 300

5 Arg Ile Leu Tyr Leu Asp Phe His Pro Val Arg Arg Gly Asp Gln Asp
 305 310 315 320

Leu Gly Tyr Val Val Thr Ile Arg Asp Arg Thr Asp Ile Ile Glu Leu
 325 330 335

10 Ser Glu Arg Leu Asp Ser Val Arg Thr Met Thr His Ala Leu Arg Ala
 340 345 350

15 Gln Arg His Glu Phe Ala Asn Arg Ile His Thr Ala Thr Gly Leu Ile
 355 360 365

Asp Ala Gly Arg Val His Asp Ala Ala Glu Phe Leu Gly Asp Ile Ser
 370 375 380

20 Arg Asn Gly Gly Gln Ser His Pro Leu Ile Gly Ser Ala His Leu Asn
 385 390 395 400

Glu Ala Phe Leu Ser Ser Phe Leu Ser Thr Ala Ser Ile Ser Ala Ser
 405 410 415

25 Glu Lys Gly Val Ser Leu Arg Ile Asn Ser Asp Thr Leu Ile Leu Gly
 420 425 430

30 Thr Val Lys Asp Pro Glu Asp Val Ala Thr Ile Leu Gly Asn Leu Ile
 435 440 445

Asn Asn Ala Ile Asp Ala Ala Val Ala Gly Glu Ala Pro Arg Trp Ile
 450 455 460

35 Glu Leu Thr Leu Met Asp Asp Ala Asp Thr Leu Val Ile Ser Val Ala
 465 470 475 480

Asp Ser Gly Pro Gly Ile Pro Glu Gly Val Asp Val Phe Ala Thr Ala
 485 490 495

40 Thr Gln Ile Gly Asp Ser Glu Asp Asn Glu Arg Thr His Gly His Gly
 500 505 510

45 Ile Gly Leu Lys Leu Cys Arg Ala Leu Ala Arg Ser His Gly Gly Asp
 515 520 525

Val Trp Val Ile Asp Arg Gly Thr Glu Asp Gly Ala Val Phe Gly Val
 530 535 540

50 Lys Leu Pro Gly Val Met Glu
 545 550

55 <210> 3
 <211> 481
 <212> DNA
 <213> Corynebacterium glutamicum

<220>

<223> citAint

<400> 3

5 ttccagtcgg tgaggtcagt gttggggttg agcgagacag tgtgtattcc cgcctgcccc 60
 tgttcctcgc cgcccttgct cttatttctg tgttgggaat ccttatcggc gtgggtgtag 120
 ccatgggcat gcgacgccgt tgggaacgcg tgaccttggg tttgcagccg gaggagctag 180
 tgacccttgt gcaaaatcag actgcagtca tcgatggcat tgatgagggc gtgctggcgc 240
 tgagcccaaa cggaacaatt ggggtgcata atgagcaggc gcaatccatg attggtgcag 300
 10 gtcctatgag tggcaggacg ttgaaagaac tagggcttga cctgggtctt gatggcgctt 360
 tattgcatgg tcagcatccg gaaaccgttg cccataacgg caggatcctc tatctggatt 420
 tccaccccggt gcgccgtggg gatcaagatt taggctacgt ggtaaccatc cgcgatcgta 480
 c 481

15

<210> 4

<211> 20

<212> DNA

<213> Corynebacterium glutamicum

20

<220>

<223> Primer citA-int1

<400> 4

25 ttccagtcgg tgaggtcagt

20

<210> 5

<211> 20

30 <212> DNA

<213> Corynebacterium glutamicum

<220>

<223> Primer citA-int2

35

<400> 5

gtacgatcgc ggatgggttac

20

[The following figure is attached:

Figure 1: Map of the plasmid pCR2.1citAint.

The abbreviations and designations used have the following meaning.

KmR:	Kanamycin resistance gene
EcoRI:	Cleavage site of the restriction enzyme EcoRI
HindIII:	Cleavage site of the restriction enzyme HindIII
BamHI:	Cleavage site of the restriction enzyme BamHI
citAint:	Internal fragment of the citA gene
ColE1:	Replication origin of the plasmid ColE1]